An $A_4 \times \mathbb{Z}_4$ model for neutrino mixing

Yoni BenTov[1], Xiao-Gang He[3,4,5], A. Zee[1,2]

[1] *Department of Physics, University of California, Santa Barbara CA 93106*
[2] *Kavli Institute for Theoretical Physics, University of California, Santa Barbara CA 93106*
[3] *INPAC, Department of Physics and Shanghai Key Laboratory for Particle Physics and Cosmology, Shanghai Jiao Tong University, Shanghai, China*
[4] *CTS, CASTS and Department of Physics, National Taiwan University, Taipei, Taiwan*
[5] *Department of Physics, National Tsing Hua University, and National Center for Theoretical Sciences, Hsinchu, 300, Taiwan*

arXiv:1208.1062v2 [hep-ph] 24 Aug 2012

Abstract

The $A_4 \times U(1)$ flavor model of He, Keum, and Volkas is extended to provide a minimal modification to tribimaximal mixing that accommodates a nonzero reactor angle $\theta_{13} \sim 0.1$. The sequestering problem is circumvented by forbidding superheavy scales and large coupling constants which would otherwise generate sizable RG flows. The model is compatible with (but does not require) a stable or metastable dark matter candidate in the form of a complex scalar field with unit charge under a discrete subgroup $\mathbb{Z}_4$ of the $U(1)$ flavor symmetry.

I Introduction

The replication of fermion generations along with small quark mixing angles and large neutrino mixing angles may suggest that the matter content of the Standard Model (SM) transforms nontrivially under a horizontal flavor symmetry group, G_F. If this flavor group G_F were a continuous global symmetry, then spontaneously breaking it would induce massless Goldstone bosons in the low-energy spectrum, which are difficult to reconcile with experiment. One might therefore imagine that either G_F is actually gauged at high energy [1] or that the global family symmetry is actually a discrete group [2] rather than a continuous Lie group.

The flavor group

$$G_F = A_4 \times U(1) \tag{I.1}$$

was originally proposed for lepton masses by E. Ma and G. Rajasekaran [3] and subsequently studied by many authors. The $U(1)$ is essentially lepton number,[1] but where heavy gauge-

[1] To avoid generating a Goldstone mode upon spontaneously breaking the $U(1)$, it is convenient to introduce a new SM-singlet scalar field charged under only a discrete subgroup $\mathbb{Z}_n$ with a self-interaction that explicitly breaks the $U(1)$. We address this issue in Section VII.

singlet antineutrinos are assigned $U(1)$ charge zero instead of -1.

In this paper we will extend a particular model by He, Keum, and Volkas (HKV) [4] which uses the same symmetry group Eq. (I.1). This model proposes that, to leading order, the CKM matrix is the identity, and the charged lepton and neutrino mass matrices are "form diagonalized" by unitary matrices that result in a tribimaximal neutrino mixing matrix,

$$V_{\rm PMNS} = V_{\rm TB} \equiv \begin{pmatrix} -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix} , \tag{I.2}$$

independently of the values of the charged lepton and neutrino masses. The main theoretical challenge for this model is that the charged fermion mass matrices are obtained using an A_4 triplet vacuum expectation value (VEV) pointing in the direction $(1,1,1)$, while the neutrino mass matrix is obtained using an A_4 triplet VEV pointing in the direction $(0,1,0)$. The former leaves a $\mathbb{Z}_3$ subgroup of A_4 unbroken, while the latter leaves a $\mathbb{Z}_2$ subgroup of A_4 unbroken. The two groups do not commute, and so a generic scalar potential will spoil the two vacuum alignments. The authors called this the "sequestering problem" and proposed one solution based on low-energy supersymmetry.

In this paper we address the sequestering problem by embedding the HKV model in a framework which has no superheavy scales, and therefore does not generate large renormalization group corrections to quadratic scalar couplings. Thus in this model the sequestering problem is solved by showing that it is not a problem to begin with: if we set various quartic couplings in the potential to be negligible at some high energy scale, they will remain small at low energy simply because the separation of scales involved is not large.

II Leading order masses and mixing

The leading order results for quark mixing and neutrino oscillations are the same as in the HKV model, so here we will only review the field transformation properties under $SU(3)\times SU(2)\times U(1)\times A_4\times U(1)_X$ and the resulting expressions for the mass matrices.

The SM fermion fields $q, \bar{u}, \bar{d}, \ell, \bar{e}$ and the SM-singlet antineutrino fields N transform as:

$$\begin{aligned} &q \sim (3,2,+\tfrac{1}{6};3,0) \ , \quad \bar{d} \sim (\bar{3},1,+\tfrac{1}{3};1+1'+1'',0) \ , \quad \bar{u} \sim (\bar{3},1,-\tfrac{2}{3};1+1'+1'',0) \\ &\ell \sim (1,2,-\tfrac{1}{2};3,+1) \ , \quad \bar{e} \sim (1,1,+1;1+1'+1'',-1) \ , \quad N \sim (1,1,0;3,0) \ . \end{aligned} \tag{II.1}$$

There are four $SU(2)\times U(1)$ Higgs doublets, $\{\Phi_a\}_{a=1}^3$ (where $a=1,2,3$ labels the triplet representation of A_4) and ϕ:

$$\Phi \sim (1,2,-\tfrac{1}{2};3,0) \ , \quad \phi \sim (1,2,-\tfrac{1}{2};1,+1) \ . \tag{II.2}$$

If we assume the A_4 VEV alignment $\langle\Phi_1^0\rangle = \langle\Phi_2^0\rangle = \langle\Phi_3^0\rangle \equiv \frac{1}{\sqrt{2}} v\, e^{i\theta_\Phi}$, the mass matrices for the down quarks, up quarks, and charged leptons are of a common form:

$$M_f = \tfrac{1}{\sqrt{2}} Y_f v\, e^{i\theta_\Phi} \ , \quad Y_f = \begin{pmatrix} y_f & y'_f & y''_f \\ y_f & \omega\, y'_f & \omega^* y''_f \\ y_f & \omega^* y'_f & \omega\, y''_f \end{pmatrix} \tag{II.3}$$

Here the index $f = d, u, e$ labels the three sectors of charged fermions ($Q = +\frac{1}{3}, -\frac{2}{3}, -1$, respectively) in the SM.

Thus the mass terms

$$\mathcal{L}_f = -(f_1, f_2, f_3) M_f \begin{pmatrix} \bar{f}_1 \\ \bar{f}_{1''} \\ \bar{f}_{1'} \end{pmatrix} + h.c. \tag{II.4}$$

are diagonalized by the unitary transformation $f = \mathcal{U}_L^f f_{\rm mass}$, where

$$\mathcal{U}_L^f = \tfrac{1}{\sqrt{3}} \begin{pmatrix} e^{\,ia_f} & e^{\,ib_f} & e^{\,ic_f} \\ e^{\,ia_f} & \omega^* e^{\,ib_f} & \omega\, e^{\,ic_f} \\ e^{\,ia_f} & \omega\, e^{\,ib_f} & \omega^* e^{\,ic_f} \end{pmatrix} \ . \tag{II.5}$$

The phase angles a, b, c are chosen to make the physical fermion masses real and positive. In Eq. (II.4) the subscripts $1, 2, 3$ label the A_4 triplet representation, while the subscripts $1, 1'', 1'$ label the different A_4 singlet representations [see Eq. (II.1)].

In this basis the Yukawa matrix is

$$(\mathcal{U}_L^f)^T Y_f = {\rm diag}(\sqrt{3}\, y_f\, e^{\,ia_f}, \sqrt{3}\, y'_f\, e^{\,ib_f}, \sqrt{3}\, y''_f\, e^{\,ic_f}) \tag{II.6}$$

so the mass of each fermion is an arbitrary coupling constant times a VEV v, just as in the SM.

The quark mixing matrix is $V_q \equiv (\mathcal{U}_L^d)^\dagger \mathcal{U}_L^u$. Putting this in the standard form $V_q \equiv \mathcal{K}_q V_{\rm CKM} \mathcal{P}_q$, we find $V_{\rm CKM} = I$. Thus this model explains why quark mixing is small; the realistic CKM angles are presumed to arise from higher order corrections, or from an extension to the theory. In this work we focus on neutrino mixing and do not pursue quark mixing any further. Consequently, from now on we suppress the $SU(3)$ quantum number when expressing group transformation properties of the fields.

As emphasized by HKV, the fermion masses are totally arbitrary while the left-handed mixing matrices are, up to unphysical phases, fixed purely by group theory, in contrast to the alternative possibility that the mixing angles might be related to ratios of masses.

III Neutrino masses

The leading contribution to neutrino masses arises through the seesaw mechanism. An A_4 triplet of SM-singlet real scalars

$$S \sim (1, 1, 0; 3, 0) \tag{III.1}$$

with an A_4 vacuum alignment $\langle S_1\rangle = \langle S_3\rangle = 0$, $\langle S_2\rangle \equiv v_S \neq 0$ (with v_S having either sign) gives a Majorana mass matrix for N:

$$M_N = m_N \begin{pmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ x & 0 & 1 \end{pmatrix} \ , \quad x \equiv \frac{y_N v_S}{m_N} \ . \tag{III.2}$$

Here m_N is a bare Majorana mass term, and y_N is the Yukawa coupling of S to $(NN)_{3_S}$. By rephasing the N fields we can take m_N real and positive without loss of generality. In this case, the parameter x is in general a complex number with an undetermined phase [4]. The Dirac mass term arises from $\mathcal{L}^\nu_{\rm Yuk} = y_\nu \phi^\dagger \ell N$ and is proportional to the identity:

$$M_D = m_D\, I \ , \quad m_D = \tfrac{1}{\sqrt{2}} y_\nu v_\phi\, e^{-i\theta_\phi} \tag{III.3}$$

where $\frac{1}{\sqrt{2}} v_\phi\, e^{i\theta_\phi} \equiv \langle \phi^0 \rangle$. In the seesaw limit the light neutrino Majorana mass matrix is

$$M_\nu \approx -M_D M_N^{-1} M_D^T = -\frac{m_D^2}{m_N} \frac{1}{1-x^2} \begin{pmatrix} 1 & 0 & -x \\ 0 & 1-x^2 & 0 \\ -x & 0 & 1 \end{pmatrix} \ . \tag{III.4}$$

This is diagonalized in the form $(\mathcal{U}^\nu_L)^T M_\nu \mathcal{U}^\nu_L = D_\nu \equiv {\rm diag}(m_1, m_2, m_3)$ by the unitary matrix

$$\mathcal{U}^\nu_L = \tfrac{1}{\sqrt{2}} \begin{pmatrix} e^{i\alpha/2} & 0 & -e^{i\gamma/2} \\ 0 & \sqrt{2}\, e^{i\beta/2} & 0 \\ e^{i\alpha/2} & 0 & +e^{i\gamma/2} \end{pmatrix} \tag{III.5}$$

where the phase angles α, β, γ are chosen such that the physical neutrino masses m_i are real and positive:

$$m_1 = \frac{m_\nu}{|1+x|} \ , \quad m_2 = m_\nu \ , \quad m_3 = \frac{m_\nu}{|1-x|} \tag{III.6}$$

where $m_\nu = |m_D^2/m_N|$. Neutrino oscillations constrain the mass-squared splittings $\Delta m^2_{\rm atm} \approx m_3^2 - m_2^2$ and $\Delta m^2_{\rm sol} \approx m_2^2 - m_1^2$ but not the overall scale m_ν. Hence to compare with experiment one computes the ratio

$$\xi \equiv \sqrt{|R|} = \begin{cases} 5.05 - 6.40 & (\Delta m^2_{\rm atm} > 0) \\ 4.93 - 6.29 & (\Delta m^2_{\rm atm} < 0) \end{cases} \qquad \text{where} \qquad R \equiv \frac{\Delta m^2_{\rm atm}}{\Delta m^2_{\rm sol}} \ . \tag{III.7}$$

This model has been shown to be compatible with Eq. (III.7) for both the "normal" mass ordering $m_1 < m_2 < m_3$ ($\Delta m^2_{\rm atm} > 0$) and the "inverted" mass ordering $m_3 < m_1 < m_2$ ($\Delta m^2_{\rm atm} < 0$) [5].

The neutrino mixing matrix is

$$V_\nu \equiv (\mathcal{U}^e_L)^\dagger \mathcal{U}^\nu_L = \begin{pmatrix} \frac{2}{\sqrt{6}}\, e^{i(\frac{\alpha}{2} - a_e)} & \frac{1}{\sqrt{3}}\, e^{i(\frac{\beta}{2} - a_e)} & 0 \\ -\frac{1}{\sqrt{6}}\, e^{i(\frac{\alpha}{2} - b_e + \frac{2\pi}{3})} & \frac{1}{\sqrt{3}}\, e^{i(\frac{\beta}{2} - b_e + \frac{2\pi}{3})} & -\frac{1}{\sqrt{2}}\, e^{i(\frac{\gamma}{2} - b_e + \frac{\pi}{6})} \\ -\frac{1}{\sqrt{6}}\, e^{i(\frac{\alpha}{2} - c_e - \frac{2\pi}{3})} & \frac{1}{\sqrt{3}}\, e^{i(\frac{\beta}{2} - c_e - \frac{2\pi}{3})} & -\frac{1}{\sqrt{2}}\, e^{i(\frac{\gamma}{2} - c_e - \frac{\pi}{6})} \end{pmatrix} \ . \tag{III.8}$$

Thus, putting this in the standard form $V_\nu \equiv \mathcal{K}_\nu V_{\rm PMNS} \mathcal{P}_\nu$ reproduces the tribimaximal mixing matrix of Eq. (I.2): $V_{\rm PMNS} = V_{\rm TB}$.

The recent measurement of a nonzero $V_{e3} \sim 0.1$ shows that this $V_{\rm PMNS}$ cannot be exact at low energy. Just as for the leading order prediction $V_{\rm CKM} \approx I$, this model predicts that $V_{\rm PMNS} \approx V_{\rm TB}$ at leading order, subject to small but nontrivial corrections.

In this work we will use the fact that the $\mathbb{Z}_3$ and $\mathbb{Z}_2$ subgroups of A_4 do not commute to generate a realistic neutrino mixing matrix. We propose a model in which the $\mathbb{Z}_2$ remains a good approximate symmetry of the neutrino sector, but corrections due to the breaking of $\mathbb{Z}_3$ feed into the neutrino mass matrix and generate the nonzero V_{e3} required to fit data.

IV Scalar potential

As explained previously, we would like the charged lepton masses to arise from a VEV $\langle \Phi^0 \rangle \propto (1,1,1)$, which leaves invariant a $\mathbb{Z}_3$ subgroup of A_4, and we would like the neutrino masses to arise from a VEV $\langle S \rangle \propto (0,1,0)$, which leaves invariant a $\mathbb{Z}_2$ subgroup of A_4. These two subgroups do not commute, and so the most general potential does not admit this configuration. This is called the sequestering problem.

In the absence of a more elaborate framework, such as extra dimensions or supersymmetry, it is unlikely that this problem can be solved completely, but it can be sufficiently mitigated in a technically natural way. The general philosophy is to declare by fiat that certain couplings are small at a particular energy scale, and then to construct a model for which these couplings do not flow substantially under the renormalization group.[2]

To make sure no superheavy scales are required, we need to bring down the scale of m_N from the usual seesaw scale; for example, if $m_N \sim$ TeV then the neutrino mass scale $m_\nu \sim 10^{-2}$ eV can be obtained if $m_D \sim m_e \sim 10^{-1}$ MeV [6]. In contrast, the simplest seesaw model has $m_N \sim M_{\rm GUT} \sim 10^{12}$ TeV and $m_D \sim v \sim 10^{-1}$ TeV. Since the Dirac mass m_D arises from the Higgs field ϕ [see Eq.(III.3)], we want to make the VEV of ϕ small while keeping its mass above the weak scale.

The problem of having heavy Higgs doublets with small VEVs was proposed and solved in the context of a different model [7]. In the present case we introduce a $G_{\rm SM} \times A_4$-invariant complex scalar field Y which carries -2 units of $U(1)_X$ charge:

$$Y \sim (1, 0; 1, -2) \tag{IV.1}$$

and a second A_4-invariant Higgs doublet

$$\phi' \sim (2, +\tfrac{1}{2}; 1, +1) \tag{IV.2}$$

[2] This is not an unfamiliar concept. In the SM, we declare by fiat that the electron Yukawa coupling is a dimensionless number of order 10^{-6}. The reason we are able to do so consistently is that a small value for this coupling is protected by chiral symmetry from receiving large corrections. In other words, we fix a small value for a dimensionless coupling and show that it does not flow significantly under RG.

with the same $U(1)_X$ charge as ϕ but with opposite hypercharge [see Eq. (II.2)]. The dimension-2 operator $\phi_i \varepsilon^{ij} \phi'_j \sim (1, 0; 1, +2)$ is invariant under $G_{\rm SM} \times A_4$ and can couple to Y to form an invariant dimension-3 operator. Consider the following scalar potential:

$$V(Y, \phi, \phi') = M_Y^2 Y^\dagger Y + M_\phi^2 \phi^\dagger \phi + M_{\phi'}^2 \phi'^\dagger \phi' + \lambda_Y (Y^\dagger Y)^2 + \lambda_{\phi'} (\phi'^\dagger \phi')^2 - (\mu Y \phi_i \varepsilon^{ij} \phi'_j + h.c.) + ... \tag{IV.3}$$

Here the ellipses contain all renormalizable terms consistent with the symmetry $G_{\rm SM} \times G_F$ whose roles are assumed subdominant in what follows. We take $M_Y^2 < 0$ and $M_{\phi'}^2 < 0$ as usual, but $M_\phi^2 > 0$. When Y and ϕ'^0 obtain VEVs, they induce a VEV for ϕ^0 according to the relation

$$v_\phi = \frac{\mu \langle Y \rangle v_{\phi'}}{M_\phi^2} + ... \tag{IV.4}$$

where $v_\phi \equiv \sqrt{2} \langle \phi^0 \rangle$ and $v_{\phi'} \equiv \sqrt{2} \langle \phi'^0 \rangle$ are assumed real and positive for simplicity. Since there is no large separation of scales in this model, it is a self-consistent (though arbitrary) assumption to declare that the contributions denoted by "..." in Eq. (IV.4), which arise from quartic interactions in the Lagrangian, are subleading. If we can drop those contributions, then $|\langle \phi^0 \rangle| \sim 1/M_\phi^2$. Thus the VEV is inversely proportional to the square of the mass. For example, if $\mu \sim v_{\phi'} \sim 10^{-1}$ TeV and $\langle Y \rangle \sim$ TeV, then

$$M_\phi \sim \left(\frac{\rm MeV}{v_\phi} \right)^{1/2} \times 10^2 \, {\rm TeV} \, . \tag{IV.5}$$

In this way we will assume that the $SU(2) \times U(1)$ doublet ϕ is the heaviest particle in the theory and thereby specify the values of all coupling constants at the scale $\mathcal{M} = M_\phi \sim 10^2$ TeV. This is much lower than the superheavy scales that often appear, e.g. $M_{\rm GUT} \sim 10^{12}$ TeV, and so the renormalization group running to scales far below $\mathcal{M}$ will not be significant, and the non-commuting $\mathbb{Z}_3$ and $\mathbb{Z}_2$ subgroups of A_4 can remain sequestered up to small corrections.

V Minimal modification to tribimaximal mixing

Consider the case for which $\mathbb{Z}_2 = \{I, r_2\}$ is unbroken but $\mathbb{Z}_3 = \{I, c, a\}$ is broken by small perturbations that feed into the neutrino mass matrix. Since ν_1 and ν_3 flip sign under $\mathbb{Z}_2$, the mass terms $\nu_1 \nu_2$ and $\nu_2 \nu_3$ are still approximately zero. However, now that $c : \nu_{1,2,3} \to \nu_{2,3,1} \to \nu_{3,1,2}$ is broken, the terms $\nu_1 \nu_1$ and $\nu_3 \nu_3$ are no longer forced to be equal. The neutrino mass matrix in the G_F-basis is now of the form [8]

$$M_\nu = \begin{pmatrix} \alpha - \varepsilon & 0 & \beta \\ 0 & \gamma & 0 \\ \beta & 0 & \alpha + \varepsilon \end{pmatrix} \, . \tag{V.1}$$

If we assume that, to a good approximation, the charged lepton mass matrix remains diagonalized by the unitary matrix studied previously[3], then the neutrino mixing matrix is a

[3]Note that this $\mathbb{Z}_2$ leaves the A_4 singlets $1, 1', 1''$ invariant. This can be seen from decomposing the product of two A_4 triplets into irreducible representations: if $u, v \sim 3$, then the three singlets are of the form

one-parameter deviation from tribimaximal mixing with the middle column $\propto (1,1,1)$ unchanged, which implies that a nonzero reactor angle θ_{13} is generated. To leading order in ε, we have

$$|V_{e3}| \approx \frac{|\varepsilon|}{\sqrt{6}|\beta|} \; . \tag{V.2}$$

To search for a suitable high-energy model that will result in the desired perturbation, we first study the low-energy effective interactions which are invariant under $G_{\rm SM} \times A_4$ between $\Phi \sim (2, -\frac{1}{2}; 3, 0)$ and the lepton doublets $\ell \sim (2, -\frac{1}{2}; 3, +1)$. Consider the dimension-5 interactions

$$\mathcal{L}_{\text{dim-5}} = c_1 \mathcal{O}_1 + c_{1'} \mathcal{O}_{1'} + c_{1''} \mathcal{O}_{1''} + c_3 \mathcal{O}_3 + h.c. \tag{V.3}$$

where:

$$\mathcal{O}_1 = (\Phi_1^\dagger \Phi_1^\dagger + \Phi_2^\dagger \Phi_2^\dagger + \Phi_3^\dagger \Phi_3^\dagger)(\ell_1 \ell_1 + \ell_2 \ell_2 + \ell_3 \ell_3) \; , \tag{V.4}$$

$$\mathcal{O}_{1'} = (\Phi_1^\dagger \Phi_1^\dagger + \omega^* \Phi_2^\dagger \Phi_2^\dagger + \omega\, \Phi_3^\dagger \Phi_3^\dagger)(\ell_1 \ell_1 + \omega\, \ell_2 \ell_2 + \omega^* \ell_3 \ell_3) \; , \tag{V.5}$$

$$\mathcal{O}_{1''} = (\Phi_1^\dagger \Phi_1^\dagger + \omega\, \Phi_2^\dagger \Phi_2^\dagger + \omega^* \Phi_3^\dagger \Phi_3^\dagger)(\ell_1 \ell_1 + \omega^* \ell_2 \ell_2 + \omega\, \ell_3 \ell_3) \; , \tag{V.6}$$

$$\mathcal{O}_3 = (\Phi_2^\dagger \Phi_3^\dagger, \Phi_3^\dagger \Phi_1^\dagger, \Phi_1^\dagger \Phi_2^\dagger) \cdot (\ell_2 \ell_3, \ell_3 \ell_1, \ell_1 \ell_2) \; . \tag{V.7}$$

These operators are invariant under $G_{\rm SM} \times A_4$ but break $U(1)_X$ by two units. If we couple any of these to a $G_{\rm SM} \times A_4$-invariant complex scalar Y with $U(1)_X$ charge -2, then these operators are promoted to dimension-6, and the coefficients c_i are proportional to $\langle Y \rangle$.

Consider the case $c_3 = 0$ and $\langle \Phi_1^0 \rangle = \langle \Phi_3^0 \rangle \neq \langle \Phi_2^0 \rangle$. Defining for convenience the overall scale of the VEVs and the small splitting as

$$a \equiv 3 \langle \Phi_1^{0\dagger} \rangle^2 \quad , \quad \delta \equiv \langle \Phi_2^{0\dagger} \rangle^2 - \langle \Phi_1^{0\dagger} \rangle^2 \tag{V.8}$$

we find:

$$\begin{aligned} c_1 \mathcal{O}_1 + c_{1'} \mathcal{O}_{1'} + c_{1''} \mathcal{O}_{1''} = {} & [a c_1 + (c_1 + \omega^* c_{1'} + \omega\, c_{1''}) \delta]\, \ell_1 \ell_1 \\ & + [a c_1 + (c_1 + c_{1'} + c_{1''}) \delta]\, \ell_2 \ell_2 \\ & + [a c_1 + (c_1 + \omega\, c_{1'} + \omega^* c_{1''}) \delta]\, \ell_3 \ell_3 \; . \end{aligned} \tag{V.9}$$

The perturbation by δ recovers a neutrino mass matrix in the form of Eq. (V.1) with:

$$\varepsilon = i\, \tfrac{\sqrt{3}}{2} (c_{1'} - c_{1''}) \delta \; . \tag{V.10}$$

Thus we seek a model in which $c_1 \sim c_{1'} \sim c_{1''} \neq 0$ and $c_3 \approx 0$, and in which $\langle \Phi^0 \rangle \propto (1,1,1)$ is perturbed in the form $\delta \langle \Phi^0 \rangle \propto (\times, 0, \times)$. In order to have the flavor-dependent[4] perturbation by δ not be overwhelmed by the flavor-independent overall shift by a [Eq. (V.8)], we could attempt to construct a model in which $c_1 \ll c_{1'} \sim c_{1''}$.

$u_1 v_1 + \omega^{p_1} u_2 v_2 + \omega^{p_2} u_3 v_3$ with $p_1, p_2 = 0, 1, 2$. Since $u_1 v_1$, $u_2 v_2$, and $u_3 v_3$ do not transform under $\mathbb{Z}_2$, neither does any linear combination of them.

[4]We are not working in the e, μ, τ basis; we use the word "flavor" here to mean that the perturbation depends on the A_4 triplet label $1, 2, 3$. This usage is consistent with the modern understanding of "flavor" as meaning "copy of representation of $G_{\rm SM}$" – in other words, copies of identical covariant derivatives – rather than having any a priori relation to the charged lepton mass basis.

VI Model for nonzero V_{e3}

One simple way to realize the effective operators from the previous section is to introduce $SU(2) \times U(1)$ triplets, $\Delta, \Delta', \Delta''$, whose transformation properties under $SU(2) \times U(1) \times A_4 \times U(1)_X$ are[5]:

$$\Delta \sim (3, -1; 1, +2) \ , \quad \Delta' \sim (3, -1; 1', +2) \ , \quad \Delta'' \sim (3, -1; 1'', +2) \ . \tag{VI.2}$$

The field $\Delta^\dagger$ can couple to $(\ell\ell)_1$, the field $\Delta'^\dagger \sim (1')^* = 1''$ to $(\ell\ell)_{1'}$, and the field $\Delta''^\dagger$ to $(\ell\ell)_{1''}$. The product $(\Delta^\dagger)^{ij}(\Phi_i\Phi_j)_1$ is invariant under $SU(2) \times U(1) \times A_4$, but it carries charge $+2$ under $U(1)_X$. Fortunately the field Y in Eq. (IV.1) has just the required property to allow the dimension-4 interactions $Y^\dagger\Delta^\dagger(\Phi\Phi)_1$, $Y^\dagger\Delta'^\dagger(\Phi\Phi)_{1'}$, and $Y^\dagger\Delta'^\dagger(\Phi\Phi)_{1''}$.

First consider the following Lagrangian for Δ:

$$\mathcal{L}_\Delta = -M_\Delta^2 \mathrm{tr}(\Delta^\dagger\Delta) + \left\{(\Delta^\dagger)^{ij}\left[\mathcal{A}(\ell_i\ell_j)_1 + \mathcal{B}^* Y^\dagger(\Phi_i\Phi_j)_1\right] + h.c.\right\} + ... \tag{VI.3}$$

Here $\mathcal{A}$ and $\mathcal{B}$ are couplings, and the ellipses denote other terms which are not important for the present discussion. We imagine that[6] $v < M_\Delta \lesssim M_\phi \sim 10^2$ TeV. At scales far below M_Δ, the path integral for Δ can be performed in the Gaussian approximation, which removes Δ from the low-energy spectrum[7] and leaves behind an effective Lagrangian:

$$\mathcal{L}_\Delta^{\text{eff}} = -\frac{1}{M_\Delta^2}\left[|\mathcal{A}|^2(\ell_i\ell_j)_1(\ell^{\dagger i}\ell^{\dagger j})_1 + |\mathcal{B}|^2|Y|^2(\Phi_i\Phi_j)_1(\Phi^{\dagger i}\Phi^{\dagger j})_1 + \left(\mathcal{A}\mathcal{B}\, Y(\Phi^{\dagger i}\Phi^{\dagger j})_1(\ell_i\ell_j)_1 + h.c.\right)\right] \ . \tag{VI.4}$$

The last term in Eq. (VI.4) is an explicit realization of the first term in Eq. (V.3), with the coefficient

$$c_1 = \mathcal{A}\mathcal{B}\,\frac{\langle Y\rangle}{M_\Delta^2} \ . \tag{VI.5}$$

This procedure can be repeated for Δ' and Δ'' to obtain the second and third operators in Eq. (V.3) with coefficients

$$c_{1'} = \mathcal{A}'\mathcal{B}'\frac{\langle Y\rangle}{M_{\Delta'}^2} \ , \quad c_{1''} = \mathcal{A}''\mathcal{B}''\frac{\langle Y\rangle}{M_{\Delta''}^2} \ . \tag{VI.6}$$

[5]We express the $SU(2)$ components of the complex triplet Δ as a 2-by-2 symmetric matrix:

$$\Delta_{ij} = \begin{pmatrix} \Delta^0 & \frac{1}{\sqrt{2}}\Delta^- \\ \frac{1}{\sqrt{2}}\Delta^- & \Delta^{--} \end{pmatrix} \tag{VI.1}$$

where the superscript denotes the electric charge.

[6]The notation "$a \lesssim b$" is to be interpreted as "a can be somewhat less than or possibly comparable to b". Also, one could instead take $M_\phi \lesssim M_\Delta$, which would amount to pushing the scale at which G_F is imposed up to $\mathcal{M} = M_\Delta$. As long as this is still substantially smaller than the usual M_{GUT}, the main results of this paper remain unchanged. See Appendix C for a toy model.

[7]Instead of working in the Wilsonian framework of performing the path integral one step at a time, the reader may prefer to study the scalar potential with all fields left in the spectrum. In that case the interaction $\Delta^\dagger Y\Phi\Phi$ of Eq. (VI.3) induces a small VEV for Δ^0, which will have the same effect as Eq. (VI.5). In the neutrino literature this is known as the Type-II seesaw mechanism [9, 10].

To leading order there is no term involving Δ, Δ', or Δ'' that couples to the A_4 triplet $(\ell_2\ell_3, \ell_3\ell_1, \ell_1\ell_2)$, so we have $c_3 = 0$. By taking M_Δ large compared to $M_{\Delta'}$ and $M_{\Delta''}$, or by taking $\mathcal{A}\mathcal{B}$ small compared to $\mathcal{A}'\mathcal{B}'$ and $\mathcal{A}''\mathcal{B}''$, we can make $c_1 \ll c_{1'}, c_{1''}$.

Now that we have generated the c_i, we need to generate the $\mathbb{Z}_3$-breaking perturbation δ. The perturbation away from $\langle\Phi^0\rangle \propto (1,1,1)$ can be achieved through a cubic interaction

$$V_{\rm cubic}(\vec{\Phi}, \vec{S}) = \tilde{\mu}_1\, \vec{S}\cdot(\vec{\Phi}^\dagger\vec{\Phi})_{3_1} + \tilde{\mu}_2\, \vec{S}\cdot(\vec{\Phi}^\dagger\vec{\Phi})_{3_2}\ . \tag{VI.7}$$

If we assume that $\langle\Phi^0_a\rangle \equiv \frac{1}{\sqrt{2}}v_a$ are real, then the two terms contribute to the classical potential through one combined invariant:

$$V_{\rm cubic}(\vec{v}, \vec{v}^S) = \tfrac{1}{2}\tilde{\mu}\,(v_1^S v_2 v_3 + v_2^S v_3 v_1 + v_3^S v_1 v_2) \tag{VI.8}$$

where $\tilde{\mu} \equiv \tilde{\mu}_1 + \tilde{\mu}_2$. This is the case studied in Appendix B. We find that $v_1 = v_3 \neq v_2$ is a solution, with

$$\delta \equiv \tfrac{1}{2}(v_1^2 - v_2^2) = \tilde{\mu}\,\frac{v_S^{(0)}}{\lambda_\Phi} + \mathcal{O}(\tilde{\mu}^2) \tag{VI.9}$$

where $v_S^{(0)}$ is the leading order value for $\langle S_2\rangle$. Using Eq. (V.2), we find[8]

$$|V_{e3}| \approx \frac{|c_{1'} - c_{1''}|}{|f(x)|}\left|\frac{\tilde{\mu}v_S^{(0)}}{2\sqrt{2}\,\lambda_\Phi m_\nu}\right| \tag{VI.10}$$

where $f(x) \equiv x/(1-x^2)$ and $x \equiv y_N v_S^{(0)}/m_N$. The various parameters in Eq. (VI.10) can be arranged to accommodate the measurement $V_{e3} \sim 0.1$.

VII Goldstone bosons and dark matter

Now that we have achieved a realistic neutrino mixing matrix, let us discuss breaking the $U(1)_X$ to a discrete subgroup to avoid generating a massless Goldstone boson after flavor symmetry breaking. Consider the $G_{\rm SM}\times A_4$-invariant scalar Y, which carries charge -2 under $U(1)_X$. Instead of the full $U(1)_X$ group, let us instead insist only on the discrete subgroup $Z_4^X : Y \to e^{i(+2)2\pi/4}Y = -Y$. Then the most general renormalizable scalar potential for Y is

$$V_Y = M_Y^2 Y^\dagger Y + (\tfrac{1}{2}m_Y^2 Y^2 + h.c.) + \lambda_Y (Y^\dagger Y)^2 + (\tfrac{1}{4}\lambda'_Y Y^4 + h.c.)\ . \tag{VII.1}$$

The terms Y^2 and Y^4 explicitly break $U(1)_X$ and remove the unwanted Goldstone mode from the low energy spectrum. The $U(1)_X$ representation $\varphi \sim m$ for the other fields should now be understood as the transformation property

$$\mathbb{Z}_4^X : \varphi \to e^{i(+m)2\pi/4}\varphi \tag{VII.2}$$

so that all fields with $m = 1$ get multiplied by a factor of i.

[8]The c_i have mass dimension -1, so V_{e3} is dimensionless as required.

Since various aspects of the model [Eq. (IV.4) and Eqs. (VI.5), (VI.6)] depend crucially on a nonzero $\langle Y \rangle$, we take the parameters in Eq. (VII.1) to be such that $\mathrm{Re}\, Y$ or $\mathrm{Im}\, Y$ (or both) obtains a nonzero VEV. There is no symmetry principle that conserves Y-number, so this SM-singlet does not provide a good candidate for dark matter.

Given this, one might be motivated to consider a $G_{\rm SM} \times A_4$-invariant complex scalar X which carries unit charge under $\mathbb{Z}_4^X$:

$$X \sim (1, 0; 1, +1) \; . \tag{VII.3}$$

The scalar potential involving only X and Y is $V(X,Y) = V_X + V_Y + V_{XY}^{\rm cubic} + V_{XY}^{\rm quartic}$, where

$$V_X = M_X^2 X^\dagger X + \lambda_X (X^\dagger X)^2 + (\tfrac{1}{4}\lambda'_X X^4 + h.c.) \; , \tag{VII.4}$$

V_Y was given in Eq. (VII.1), and:

$$V_{XY}^{\rm cubic} = \left[\tfrac{1}{2}(\rho\, Y + \rho'\, Y^\dagger) X^2 + h.c.\right] \; , \tag{VII.5}$$

$$V_{XY}^{\rm quartic} = \lambda_{XY} X^\dagger X\, Y^\dagger Y + \left(\tfrac{1}{2}\lambda'_{XY} X^\dagger X\, Y^2 + h.c.\right) \; . \tag{VII.6}$$

If we take M_X^2 positive so that V_X does not induce a VEV for X, then even when Y picks a VEV and breaks $\mathbb{Z}_4^X$, the potential $V(X,Y)$ leaves a discrete X-parity $P_X : X \to -X$ unbroken. Thus, if the (X,Y)-sector of the theory were secluded from the rest of the Lagrangian, the lighter of $\mathrm{Re}(X)$ and $\mathrm{Im}(X)$ would be stable and provide a candidate for dark matter [11, 12].

Now consider the interaction of X with the other fields. In addition to the obvious "Higgs-portal" interactions such as $X^\dagger X \Phi^\dagger \Phi$ which conserve X-number, there are quartic interactions which are linear in X:

$$V_{\not P_X} = \kappa_1 X \phi^{\dagger i} \vec{\Phi}_i \cdot \vec{S} + \kappa_2 X \phi'^{\dagger i} \varepsilon_{ij} \vec{\Phi}^{\dagger j} \cdot \vec{S} + h.c. \tag{VII.7}$$

If we suppose that κ_1 or κ_2 is small but nonzero, then P_X is broken and X can decay. In view of the scales in our model, one might suppose that $M_X \gtrsim$ TeV so that X decays either to the three-body final state $\phi^* \Phi S$ or $\phi' \Phi S$, or if such a decay is kinematically inaccessible, then X can decay to whatever final states can be reached through the propagators of virtual scalars, e.g. TeV-scale neutrinos or SM charged leptons. For the sake of a rough calculation suppose that $M_X \gg M_{\phi'} + M_\Phi + M_S$ and estimate the width (inverse lifetime) of X as

$$\tau_X^{-1} \equiv \Gamma_X \sim \frac{\kappa^2}{(2\pi)^n} M_X \tag{VII.8}$$

where κ is either κ_1 or κ_2, and $(2\pi)^n$ is a phase space factor for n outgoing particles. Recent measurements of cosmic ray spectra [13, 14] may indicate the presence of late-decaying dark matter, which would require [15] a lifetime much larger than the age of the universe: $\tau_{\rm min} \sim 10^{27}\mathrm{s} \sim 10^{54}\ \mathrm{TeV}^{-1}$, where as usual we work in units $\hbar = 1$. Thus, if we would like to interpret the cosmic ray anomalies in terms of our metastable X particle, we need:

$$\frac{\tau_X}{\tau_{\rm min}} \sim (2\pi)^n \left(\frac{\mathrm{TeV}}{M_X}\right) \left(\frac{10^{-27}}{\kappa}\right)^2 \gtrsim 1 \; . \tag{VII.9}$$

Thus, as is the usual case with a decay rate resulting from renormalizable interactions, we require extremely small coupling constants, $\kappa \lesssim 10^{-27}$, if we insist on a DM interpretation of the data. It is worth recalling [see below Eq. (VII.6)] that if we were to impose $P_X : X \to -X$ on the model, then we would have $\kappa_1 = \kappa_2 = 0$. Taking this argument in the other direction, we note that in the limit $\kappa_{1,2} \to 0$ the transformation $P_X : X \to -X$ emerges as a symmetry of the Lagrangian. Thus any value of κ in Eq. (VII.9), no matter how small, is technically natural in the sense of 't Hooft. In order to satisfy Eq. (VII.9) we would need to fix κ_1 and κ_2 to be small dimensionless numbers, but we would not need to engineer any unnatural cancellation between large numbers.

VIII Discussion

We have embedded the HKV model into a framework whose largest scale $\mathcal{M}$ is much smaller than the traditional seesaw scale: $\mathcal{M}/M_{\rm GUT} \sim 10^{-10}$. The main requirements of such a scenario are an additional A_4-singlet Higgs doublet, ϕ', and an SM-singlet complex scalar, Y, which carry $U(1)_X$ charge $+1$ and -2, respectively [Eqs. (IV.1) and (IV.2)]. Then the scalar potential can be arranged such that the small Dirac mass for the neutrino can be generated by a Higgs doublet whose VEV is small but whose mass is large [Eq. (IV.4)].

This ameliorates the sequestering problem that typically afflicts models of this type for the simple reason that the separation between the scale at which $G_F = A_4 \times U(1)_X$ is imposed and the scale of neutrino masses is not sufficiently vast to induce large renormalization group corrections. This is an alternative to the supersymmetric and brane-world approaches mentioned by HKV and studied by various authors.

The recent measurement of $V_{e3} \sim 0.1$ is accommodated by a minimal modification to tribimaximal mixing due to A_4-singlet Higgs triplets which communicate the breaking of $\mathbb{Z}_3 = \{I, c, a\}$ into the neutrino sector while leaving the $\mathbb{Z}_2 = \{I, r_2\}$ essentially unbroken (see Section VI).

The desire to remove an unwanted Goldstone boson due to breaking the continuous global $U(1)_X$ led us to insist only on the discrete subgroup $\mathbb{Z}_4^X < U(1)_X$, and the Lagrangian for the SM-singlet Y explicitly implements the distinction between the two symmetry groups [Eq. (VII.1)]. This then motivated (but did not require) the introduction of a second SM-singlet complex scalar field X with half the $\mathbb{Z}_4^X$ charge of Y, which can be a candidate for decaying dark matter [Eqs. (VII.3) and (VII.7)].

We conclude by mentioning other possibilities for neutrino mixing in this framework; the one-parameter deviation from tribimaximal mixing studied here is by no means the only option [16]. For example, the study of Appendix B might lead us to consider the case where $\mathbb{Z}_3 = \{I, c, a\}$ remains a good symmetry of the neutrino mass matrix, while the breaking of $\mathbb{Z}_2 = \{I, r_2\}$ becomes significant. In this case, the neutrino mass matrix in the G_F-basis

becomes

$$M_\nu = \begin{pmatrix} \alpha & \varepsilon_{12} & \beta \\ \varepsilon_{12} & \gamma & \varepsilon_{23} \\ \beta & \varepsilon_{23} & \alpha \end{pmatrix} \tag{VIII.1}$$

with small parameters ε_{ij}. If $\mathcal{U}_L^e$ remains approximately unchanged, then $V_{e3} = [(\mathcal{U}_L^e)^\dagger \mathcal{U}_L^\nu]_{e3} \neq 0$ must arise by modifying the third column of $\mathcal{U}_L^\nu$. For example,

$$M_\nu \begin{pmatrix} \pm 1 \\ 0 \\ 1 \end{pmatrix} = (\alpha \pm \beta) \begin{pmatrix} \pm 1 \\ 0 \\ 1 \end{pmatrix} + (\varepsilon_{12} \pm \varepsilon_{23}) \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \ . \tag{VIII.2}$$

Note that if $\varepsilon_{12} = +\varepsilon_{23}$, then the third column of $\mathcal{U}_L^\nu$ remains unchanged from its leading order value, which is not what we want.

Acknowledgments:

This research was supported by the NSF under Grant No. PHY07-57035.

A A_4 invariant potential for a real triplet

The scalar potential for a Higgs field, φ, transforming as an A_4 triplet was studied previously by one of the authors [17]. Invariance under $SU(2) \times U(1)$ implies that this potential is a function only of the combination $\varphi_a^\dagger \varphi_b$, where a, b label the A_4 components, and it has one quadratic invariant and five quartic invariants.

Here we study the scalar potential for a real scalar field S transforming as a triplet under A_4. Since S is real, the five quartic invariants for φ collapse to only two independent terms. However, since we do not impose a parity that flips the sign of S (the real-valued analog of a $U(1)$ symmetry), there is also a nontrivial cubic invariant from the group theoretic property $3 \times 3 \times 3 = 3 \times (1 + 1' + 1'' + 3_A + 3_S) = 3 \times 3_S + ... = 1 + ...$, where the 1 denotes the invariant $S_1 S_2 S_3$. (Note that $3_A = 0$ since $S_a S_b = S_b S_a$.)

The most general renormalizable, A_4-invariant scalar potential for $S \sim 3$ is

$$V(S) = \tfrac{1}{2} M^2 \sum_{a=1}^{3} S_a^2 + \mu S_1 S_2 S_3 + \tfrac{1}{4} \lambda \left(\sum_{a=1}^{3} S_a^2 \right)^2 + \tfrac{1}{2} \lambda' \left[(S_1 S_2)^2 + (S_2 S_3)^2 + (S_3 S_1)^2 \right] \ . \tag{A.1}$$

We study three[9] potential vacua of interest: $E : \{\langle S_1 \rangle = \langle S_2 \rangle = \langle S_3 \rangle \equiv S \neq 0\}$, $U : \{\langle S_1 \rangle \equiv S \neq 0, \langle S_2 \rangle = \langle S_3 \rangle = 0\}$, and $P : \{\langle S_1 \rangle = \langle S_2 \rangle \equiv S \neq 0, \langle S_3 \rangle = 0\}$.

[9]The case $\langle S_2 \rangle = \langle S_3 \rangle = -\langle S_1 \rangle$ was studied by E. Ma in the context of another model based on A_4 symmetry [18].

In the vacuum E we have $M^2 = -(3\lambda + 2\lambda')S^2 - \mu S$, and:

$$V|_E = -\tfrac{9}{4}(\lambda + \tfrac{2}{3}\lambda')S^4 - \tfrac{1}{2}\mu S^3 \qquad \text{and} \qquad \left.\frac{\partial^2 V}{\partial S_a \partial S_b}\right|_E = \begin{pmatrix} m^2 & \Delta & \Delta \\ \Delta & m^2 & \Delta \\ \Delta & \Delta & m^2 \end{pmatrix} \tag{A.2}$$

where $m^2 \equiv 2\lambda S^2 - \mu S$, $\Delta \equiv 2(\lambda + \lambda')S^2 + \mu S$. The notation $V|_E$ denotes the value of the potential evaluated at the minimum for the vacuum E. Diagonalizing the matrix of second derivatives gives $\mathcal{D}^2_E = \mathrm{diag}(m^2 + 2\Delta, m^2 - \Delta, m^2 - \Delta)$. The requirement $m^2 - \Delta \geq 0$ implies $\mu S \leq -\frac{1}{2}\lambda' S^2$.

In the vacuum U we have $M^2 = -\lambda S^2$, and:

$$V|_U = -\tfrac{1}{4}\lambda S^4 \qquad \text{and} \qquad \left.\frac{\partial^2 V}{\partial S_a \partial S_b}\right|_U = \begin{pmatrix} m_0^2 & 0 & 0 \\ 0 & m^2 & \Delta \\ 0 & \Delta & m^2 \end{pmatrix} \tag{A.3}$$

where $m_0^2 \equiv 2\lambda S^2$, $m^2 \equiv (\lambda' - \lambda)S^2$, and $\Delta \equiv \mu S$. The matrix of mass-squared eigenvalues is $\mathcal{D}^2_U = \mathrm{diag}(m_0^2, m^2 - 2\Delta, m^2 + 2\Delta)$. The requirement $m^2 - 2\Delta \geq 0$ is satisfied if $\mu S \leq \frac{1}{2}(\lambda' - \lambda)S^2$.

When studying the putative vacuum P we find an important difference from the previously studied Higgs potential. Minimizing the potential with respect to S_3 results in the equation

$$\left.\frac{\partial V}{\partial S_3}\right|_P = \mu S^2 = 0 \; . \tag{A.4}$$

If $\mu \neq 0$ then this equation cannot be satisfied. Thus, unlike the case of a complex $SU(2) \times U(1)$-invariant Higgs potential, a real scalar triplet does not admit the vacuum P even as a local minimum.[10]

B Interactions between a Higgs field and a real triplet

Now we study in detail whether the VEVs $\langle S \rangle \propto (1,0,0)$ and $\langle \varphi \rangle \propto (1,1,1)$ can hold to a good approximation in a Lagrangian based on $A_4 \times \mathbb{Z}_2$. Let $\varphi \sim 3_-$ and $S \sim 3_+$ be two real triplets of A_4, where the subscript indicates the parity under $\mathbb{Z}_2$. The real field φ which is odd under $\mathbb{Z}_2$ serves as a toy model for a complex field Φ which is charged under $SU(2) \times U(1)$.

The potential is $V = V_S + V_\varphi + V_{S\varphi}^{\text{cubic}} + V_{S\varphi}^{\text{quartic}}$, where

$$V_S = \tfrac{1}{2}M_S^2 \sum_{a=1}^{3} S_a^2 + \mu_S S_1 S_2 S_3 + \tfrac{1}{4}\lambda_S \left(\sum_{a=1}^{3} S_a^2\right)^2 + \tfrac{1}{2}\lambda'_S \left[(S_1 S_2)^2 + (S_2 S_3)^2 + (S_3 S_1)^2\right] \tag{B.1}$$

$$V_\varphi = \tfrac{1}{2}M_\varphi^2 \sum_{a=1}^{3} \varphi_a^2 + \tfrac{1}{4}\lambda_\varphi \left(\sum_{a=1}^{3} \varphi_a^2\right)^2 + \tfrac{1}{2}\lambda'_\varphi \left[(\varphi_1 \varphi_2)^2 + (\varphi_2 \varphi_3)^2 + (\varphi_3 \varphi_1)^2\right] \tag{B.2}$$

[10] In the $SU(2) \times U(1)$ case P is a local minimum but never the global one [17].

describes each sector, and

$$V_{S\varphi}^{\text{cubic}} = \tilde{\mu}(S_1\varphi_2\varphi_3 + S_2\varphi_3\varphi_1 + S_3\varphi_1\varphi_2) \tag{B.3}$$

$$V_{S\varphi}^{\text{quartic}} = \kappa_0(\sum_a S_a^2)(\sum_b \varphi_b^2) + \kappa_1(S_1^2\varphi_2^2 + \text{cyclic}) + \kappa_2(\varphi_1^2 S_2^2 + \text{cyclic}) + \kappa_3(S_1 S_2 \varphi_1\varphi_2 + \text{cyclic}) \tag{B.4}$$

describe the interactions between the two sectors.

We would like to find a perturbative solution around the configurations $S \propto (1,0,0)$ and $\varphi \propto (1,1,1)$. The general case can be parametrized as

$$S_a = S_a^{(0)} + \varepsilon S_a^{(1)} \ , \quad \varphi_a = \varphi_a^{(0)} + \varepsilon\varphi_a^{(1)} \tag{B.5}$$

where ε is a small parameter. We will first assume that $V_{S\varphi}^{\text{quartic}}$ can be dropped, or in other words that the κ_i are at most $\mathcal{O}(\varepsilon^2)$ and thus negligible. We will assume that $V_{S\varphi}^{\text{cubic}} \equiv \varepsilon V_1$ is itself a perturbation by defining $\tilde{\mu} = \varepsilon\tilde{\mu}_0$ with $\tilde{\mu}_0$ a parameter of $\mathcal{O}(\mu_S)$. The potential $V_0 \equiv V_S + V_\varphi$ is to be treated as the leading order potential of $\mathcal{O}(\varepsilon^0)$, with any possible renormalization group corrections to its parameters being of $\mathcal{O}(\varepsilon^2)$ or higher.

The goal is to solve the equations $\frac{\partial}{\partial S_a}(V_0 + \varepsilon V_1) = 0$ and $\frac{\partial}{\partial \varphi_a}(V_0 + \varepsilon V_1) = 0$ to $\mathcal{O}(\varepsilon)$. In the particular leading order vacuum of interest, $S_a^{(0)} \propto (1,0,0)$ and $\varphi_a^{(0)} \propto (1,1,1)$, this procedure gives the result

$$S_a = \begin{pmatrix}1\\0\\0\end{pmatrix}_a S_0 + \varepsilon S_a^{(1)} \ , \quad \varphi_a = \begin{pmatrix}1\\1\\1\end{pmatrix}_a \varphi_0 + \varepsilon\varphi_a^{(1)} \tag{B.6}$$

where:

$$S_1^{(1)} = -\frac{1}{2\lambda_S}\left(\frac{\varphi_0}{S_0}\right)^2 \tilde{\mu}_0 \ , \quad S_2^{(1)} = S_3^{(1)} = -\frac{1}{\lambda'_S + \frac{\mu_S}{S_0}}\left(\frac{\varphi_0}{S_0}\right)^2 \tilde{\mu}_0 \ , \tag{B.7}$$

$$\varphi_1^{(1)} = -\frac{\lambda_\varphi + \lambda'_\varphi}{\lambda_\varphi(\lambda_\varphi + \frac{3}{2}\lambda'_\varphi)}\left(\frac{S_0}{\varphi_0}\right)\tilde{\mu}_0 \ , \quad \varphi_2^{(1)} = \varphi_3^{(1)} = +\frac{\lambda'_\varphi}{2\lambda_\varphi(\lambda_\varphi + \frac{3}{2}\lambda'_\varphi)}\left(\frac{S_0}{\varphi_0}\right)\tilde{\mu}_0 \ . \tag{B.8}$$

If we take $\varphi_0/S_0 \lesssim 1$, then the corrections to $S_a \propto (1,0,0)$ can be ignored safely, but those to $\varphi_a \propto (1,1,1)$ are important. This is precisely the situation we seek in the main text: the $\mathbb{Z}_3$ "charged lepton" subgroup of A_4 is broken, but the $\mathbb{Z}_2$ "neutrino" subgroup of A_4 remains conserved to a good approximation. The effect we are talking about is of order $S_2^{(1)}/(\varphi_2^{(1)} - \varphi_1^{(1)}) \sim (\varphi_0/S_0)^3$, e.g. $\sim (1/4)^3 \sim 1\%$.

A convenient way to parametrize this breaking of $\mathbb{Z}_3 < A_4$ is:

$$\delta \equiv \tfrac{1}{2}(\varphi_2^2 - \varphi_1^2) = \varepsilon\tilde{\mu}_0\frac{S_0}{\lambda_\varphi} + \mathcal{O}(\varepsilon^2) \ . \tag{B.9}$$

This is the equation we use to obtain Eq. (VI.9) in the main text.

C RG for toy model coupled to fermions

Consider a simple toy model of two real scalars ϕ_1, ϕ_2 coupled to "quarks" q and "antiquarks" $\bar{q}$, with a color gauge group $SU(N)$ under which q and $\bar{q}$ transform as N and N^*, respectively. (The ϕ_i are color-neutral, and the gluons are denoted by $\{G_\mu^a\}_{a=1}^{N^2-1}$.) We impose the reflection symmetry[11]

$$\mathbb{Z}_2 : \qquad \phi_1 \to -\phi_1,\ q \to -q,\ \phi_2 \to +\phi_2,\ \bar{q} \to +\bar{q} \tag{C.1}$$

so that ϕ_1 has a Yukawa interaction with $q\bar{q}$ whereas ϕ_2 does not:

$$\mathcal{L}_{\text{Yuk}} = -y\phi_1 q\bar{q} + h.c. \tag{C.2}$$

The most general renormalizable potential for ϕ_1 and ϕ_2 consistent with $\mathbb{Z}_2$ symmetry is:

$$V(\phi_1,\phi_2) = \tfrac{1}{2}M_1^2\phi_1^2 + \tfrac{1}{2}M_2^2\phi_2^2 + M_{12}\,\phi_1^2\phi_2 + \tfrac{1}{4!}\lambda_1\phi_1^4 + \tfrac{1}{4!}\lambda_2\phi_2^4 + \tfrac{1}{4}\lambda_{12}\phi_1^2\phi_2^2\ . \tag{C.3}$$

We are interested in the renormalization group equations (RGE) for the Lagrangian

$$\mathcal{L}_{\text{toy}}(G,\phi_1,\phi_2,q,\bar{q}) = \mathcal{L}_{\text{kin}}(G,\phi_1,\phi_2,q,\bar{q}) + \mathcal{L}_{\text{Yuk}}(\phi_1,q,\bar{q}) - V(\phi_1,\phi_2) \tag{C.4}$$

in the deep UV regime, far above the scalar masses M_1, M_2 and the dimensionful cubic coupling M_{12}. (Since we are not interested in low-energy phenomena such as spontaneous symmetry breaking, we take M_1^2 and M_2^2 both positive.)

The issue we would like to address is the following. Let m denote a "low-energy" mass scale $m \sim M_1 \sim M_2 \sim M_{12}$, and let $\mathcal{M}$ denote the UV cutoff of the field theory. Suppose we (arbitrarily) fix the parameters of the theory at the scale $\mathcal{M}$ to satisfy

$$\lambda_1(\mathcal{M}) \sim \lambda_2(\mathcal{M}) \sim O(1)\ , \quad |\lambda_{12}(\mathcal{M})| \ll 1\ . \tag{C.5}$$

Our goal is then to find the maximum ratio $\mathcal{M}/m$ such that the "off-diagonal" coupling λ_{12} remains parametrically smaller than the "diagonal" couplings λ_1 and λ_2 at energy $\mu \sim m \ll \mathcal{M}$.

This toy model is a simple subcase of a general class of theories studied by Cheng, Eichten, and Li [19]. Following their notation, we express our Lagrangian as[12]

$$\mathcal{L} = -\tfrac{1}{4}G^a_{\mu\nu}G^{a\mu\nu} - \tfrac{1}{2}\partial_\mu\phi_i\partial^\mu\phi_i + iq^\dagger\bar{\not{D}}q + i\bar{q}^\dagger\bar{\not{D}}\bar{q} - (q\,h_i\bar{q}\,\phi_i + h.c.) - V(\phi)\ , \tag{C.6}$$

$$V(\phi) = \tfrac{1}{4!}f_{ijk\ell}\phi_i\phi_j\phi_k\phi_\ell + \text{lower-dimension terms}\ . \tag{C.7}$$

Here $G^a_{\mu\nu} = \partial_\mu G^a_\nu - \partial_\nu G^a_\mu + gC^{abc}G^b_\mu G^c_\nu$ is the gluon field strength, and $\bar{\not{D}} = \bar{\sigma}^\mu D_\mu$ is the covariant derivative for each 2-component fermion: $D_\mu q = (\partial_\mu - igG^a_\mu t^a)q$ and $D_\mu\bar{q} = (\partial_\mu - igG^a_\mu\bar{t}^a)\bar{q}$ with $\bar{t}^a = -(t^a)^*$.

[11] This $\mathbb{Z}_2$ serves as a toy model for $SU(2)$ gauge symmetry, under which Higgs fields ("ϕ_1") and left-handed quarks transform, but under which SM-singlet scalars ("ϕ_2") and left-handed antiquarks are neutral.

[12] In contrast to the more general case studied in the reference, but similarly to the realistic case in the SM and many of its extensions, here our scalars ϕ_i are not charged under the color group. Also here we need only one flavor of quark, in which case h_i is a single number (for each i) rather than a matrix.

The lowest-order[13] RGE for the couplings g, h_i, and $f_{ijk\ell}$ are ($\mathcal{D} \equiv 16\pi^2 d/dt$, $t \equiv \ln \frac{\mu}{\mu_0}$ for the scale μ and an arbitrary reference point μ_0):

$$\mathcal{D}g = -\tfrac{1}{2} b_0 g^3 \ , \quad b_0 = 2\left(\tfrac{11}{3} S_1^{\mathcal{G}} - \tfrac{4}{3} S_3^F\right) \tag{C.8}$$

$$\begin{aligned}\mathcal{D}h_i =& 2h_m h_i h_m + \tfrac{1}{2}(h_m h_m h_i + h_i h_m h_m) + 2\mathrm{tr}(h_i h_m) h_m \\ & -3(2t^a h_i t^a + S_2^F h_i) g^2 + \tfrac{1}{288\pi^2} f_{ijk\ell} f_{jk\ell m} h_m\end{aligned} \tag{C.9}$$

$$\begin{aligned}\mathcal{D}f_{ijk\ell} =& f_{ijmn} f_{mnk\ell} + f_{ikmn} f_{mnj\ell} + f_{i\ell mn} f_{mnjk} + 8\mathrm{tr}(h_i h_m) f_{mjk\ell} \\ & -2\mathrm{tr}(h_i h_j \{h_k, h_\ell\} + h_i h_k \{h_j, h_\ell\} + h_i h_\ell \{h_j, h_k\})\end{aligned} \tag{C.10}$$

Here we have defined the group theoretic quantities $C^{acd} C^{bcd} = S_1^{\mathcal{G}} \delta^{ab}$, $t^a t^a = S_2^F I$, and $\mathrm{tr}(t^a t^b) = S_3^F \delta^{ab}$ related to the color representation of the fermions[14]. For $SU(N)$, we have $S_1^{\mathcal{G}} = N$, $S_2^F = \frac{N^2-1}{2N}$, and $S_3^F = \frac{1}{2}$.

For our case [Eqs. (C.2) and (C.3)] we have:

$$h_i = y\, \delta_{i1} \ , \tag{C.11}$$

$$f_{1111} = \lambda_1 \ , \quad f_{2222} = \lambda_2 \ , \tag{C.12}$$

$$f_{1122} = f_{1212} = f_{1221} = f_{2121} = f_{2112} = f_{2211} = \lambda_{12} \ , \tag{C.13}$$

$$f_{1112} = f_{1121} = f_{1211} = f_{2111} = f_{2221} = f_{2212} = f_{2122} = f_{2221} = 0 \ . \tag{C.14}$$

Then Eq. (C.9) simplifies to

$$\mathcal{D}y = 5y^3 - 9 S_2^F y g^2 + \tfrac{1}{288\pi^2}(\lambda_1^2 + 3\lambda_{12}^2) y \tag{C.15}$$

and Eq. (C.10) produces three reasonably simple coupled equations:

$$\mathcal{D}\lambda_1 = 3(\lambda_1^2 + \lambda_{12}^2) + 8y^2 \lambda_1 - 12 y^4 \tag{C.16}$$

$$\mathcal{D}\lambda_2 = 3(\lambda_2^2 + \lambda_{12}^2) \tag{C.17}$$

$$\mathcal{D}\lambda_{12} = (\lambda_1 + \lambda_2)\lambda_{12} + 4\lambda_{12}^2 + 8y^2 \lambda_{12} \tag{C.18}$$

Before solving the system numerically, let us first obtain some intuition for how the "off-diagonal" coupling λ_{12} flows. The whole point of our study is to consider the case for which λ_{12} is parametrically smaller than the other couplings, so suppose we can drop terms of $O(\lambda_{12}^2)$. If we drop the $O(\lambda_{12}^2)$ term in Eq. (C.18), then the flow equation for λ_{12} is of the form $\mathcal{D}\lambda_{12} = \mathcal{F}(\mu)\lambda_{12}(\mu)$, with $\mathcal{F} = \lambda_1 + \lambda_2 + 4\lambda_{12} + 8y^2$, and the equation can be integrated:

$$\lambda_{12}(m) \approx \lambda_{12}(\mathcal{M})\, e^{-\int_m^{\mathcal{M}} \frac{d\mu}{\mu} \mathcal{F}(\mu)} \ . \tag{C.19}$$

[13] As pointed out in the reference, the last term of Eq. (C.9) is of "lowest-order" even though it arises from a 2-loop diagram.

[14] Our two-component quarks are coupled in a "vectorlike" fashion to the gluons, so it is convenient to package them into 4-component Dirac spinors $Q = (q, \bar{q}^\dagger)$, which here transform under the defining N-dimensional representation of $SU(N)$.]

Thus in the limit $\lambda_{12}(\mathcal{M}) \to 0$, the coupling $\lambda_{12}(\mu)$ gets renormalized multiplicatively, so perhaps contrary to intuition, we should expect that the assumption $\lambda_{12}(\mu) \ll 1$ remains valid even for $\mu \sim m \ll \mathcal{M}$.

Now we solve the system of ordinary differential equations {(C.8), (C.15)-(C.18)} numerically. For the high energy scale we take the extreme case $\mathcal{M} \approx 10^{15}$ GeV and we flow to the weak scale, $m \approx 10^2$ GeV, in which case $t \equiv \ln(\mu/\mathcal{M})$ takes values approximately in the range $0 \geq t \geq -30$.

In the SM the strong coupling (in the $\overline{\text{MS}}$ renormalization scheme for concreteness) at the Z pole mass is known to be [20] $g_S(M_Z) \equiv \sqrt{4\pi\alpha_S(M_Z)} \approx 1.22$ and runs to ~ 0.5 at the scale of grand unification in a typical $SU(5)$ model with a desert between the two scales. Thus for illustrative purposes we take the initial condition $g(\mathcal{M}) = 0.5$.

The Yukawa coupling in our toy model should be thought of as the case in which the top quark couples to the "Higgs-like" scalar ϕ_1. In the model studied in the main text, the A_4-triplet Higgs field Φ is assumed to give mass to both the charged leptons and the quarks, so the situation $y \sim y_t \sim 1$ is the one appropriate for the present study. Thus we take $y(\mathcal{M}) = 1$.

For the scalar interactions, the purpose is to study the case $\lambda_1(\mathcal{M}) \sim \lambda_2(\mathcal{M}) \sim 1$ and[15] $|\lambda_{12}(\mathcal{M})| \ll 1$. To be concrete we take $\lambda_1(\mathcal{M}) = \lambda_2(\mathcal{M}) = 1$ and $\lambda_{12}(\mathcal{M}) = 10^{-3}$.

The running couplings $g(\mu), y(\mu), \lambda_1(\mu), \lambda_2(\mu)$, and $\lambda_{12}(\mu)$ are displayed in Figs. 1(a), 1(b), 2(a), 2(b), and 3, respectively. The gauge coupling blows up in the deep infrared, as expected, and carries the Yukawa coupling with it into the nonperturbative regime. Both couplings remain ~ 1 down to $t \sim -25$, or $\mu/\mathcal{M} = e^t \sim 10^{-11}$.

The quartic self-coupling λ_1 remains essentially fixed to its high-scale value until about the same scale at which g and y become non-perturbative, and the quartic-self coupling λ_2 runs down slowly from $\lambda_2(\mathcal{M}) = 1$ to $\lambda_2(e^{-25}\mathcal{M}) \sim 0.6$. The difference in running between the two is of course due to the different roles of ϕ_1 and ϕ_2 in the model: the gauge coupling feeds into the running for the Yukawa coupling, which contributes only to interactions involving ϕ_1 and not ϕ_2.

The main figure of interest is Fig. 3, which shows that the off-diagonal coupling λ_{12} runs down from its high-scale value $\lambda_{12}(\mathcal{M}) = 10^{-3}$ to ever smaller values, down to $\lambda_{12}(e^{-25}\mathcal{M}) \sim 2\times10^{-4}$. Extrapolating these qualitative results to the model in the paper, we expect that the

[15] In a two-Higgs doublet model with Higgs doublets ϕ_1 and ϕ_2, the VEVs must align in the charge-conserving vacuum so that the photon does not obtain a mass. This requires the coefficient of the interaction $|\phi_1^\dagger\phi_2|^2$ to be negative. Furthermore, if $\text{Im}(\langle\phi_1\rangle\langle\phi_2\rangle) = 0$ then the coefficient of $(\phi_1^\dagger\phi_2)^2 + h.c.$ should also be negative. [21] In the present case, the coupling λ_{12} stands roughly for these two interactions as well as for the coefficient of $|\phi_1|^2|\phi_2|^2$. The running of λ_{12} will be slow enough such that its sign does not flip, and so if we extrapolate to the $SU(2)\times U(1)$-invariant case we can just pick the appropriate sign at the high energy scale $\mathcal{M}$, which is taken much less than the typical GUT scale, such that electromagnetism remains unbroken at low energy.

rather mild separation of scales between the weak scale and the "heavy" mass scales of either the Higgs doublet ϕ [Eqs. (II.2), (IV.4), (IV.5)] or the Higgs triplet Δ [Eqs. (VI.2), (VI.5)] does not spoil the small initial values assumed for the quartic self-couplings in the scalar potential, which would mix the "charged lepton" and "neutrino" sectors of the model and introduce the sequestering problem.

References

[1] H. Georgi, "Towards a grand unified theory of flavor," Nucl. Phys. B, Vol. 156, No. 1, 27 Aug 1979, pp. 126-134

[2] A. de Rújula, H. Georgi, and S. L. Glashow, "A theory of flavor mixing," Ann. Phys. 109, 258-313 (1977) ; F. Wilczek and A. Zee, "Discrete flavor symmetries and a formula for the Cabibbo angle," Phys. Lett. B, Vol. 70, No. 4, 24 Oct 1977, pp. 418-420; G. Sartori, "Discrete symmetries, natural flavor conservation and weak mixing angles," Phys. Lett. B, Vol. 82, No. 2, 26 Mar 1979, pp. 255-259; E. Derman and H. S. Tsao, "$[SU(2)\times U(1)]\times S_n$ flavor dynamics and a bound on the number of flavors," Phys. Rev. D, Vol. 20, No. 5, 1 Sep 1979, pp. 1207-1215

[3] E. Ma and G. Rajasekaran, "Softly broken A_4 symmetry for nearly degenerate neutrino masses," Phys. Rev. D, Vol. 64, 113012 (2001)

[4] X. G. He, Y. Y. Keum, and R. R. Volkas, "A_4 flavour symmetry breaking scheme for understanding quark and neutrino mixing angles," JHEP 0604:039,2006 (arXiv:hep-ph/0601001v4)

[5] K. S. Babu and X. G. He, "Model of Geometric Neutrino Mixing," arXiv:hep-ph/0507217v2

[6] W. Grimus and L. Lavoura, "TeV-scale seesaw mechanism catalyzed by the electron mass," Phys.Lett.B687:188-193,2010 (arXiv:0912.4361v3 [hep-ph])

[7] R. A. Porto and A. Zee, "The private Higgs," Phys.Lett.B666:491-495,2008 (arXiv:0712.0448v3 [hep-ph])

[8] X. G. He and A. Zee, "Minimal modification to the tri-bimaximal neutrino mixing," Phys. Lett. B645 (2007) 427-431; X. G. He and A. Zee, "Minimal modification to tribimaximal mixing," Phys. Rev. D 84, 053004 (2011)

[9] J. Schechter and J. W. F. Valle, "Neutrino masses in $SU(2)\times U(1)$ theories," Phys. Rev. D, Vol. 22, No. 9, 1 Nov. 1980, pp. 2227 - 2235.

[10] G. Lazarides and Q. Shafi, "Proton lifetime and fermion masses in an $SO(10)$ model," Nucl. Phys. B, Vol. 181, No. 2, 6 Apr 1981, pp. 287-300; R. N. Mohapatra and G. Senjanović, "Neutrino masses and mixings in gauge models with spontaneous parity violation," Phys. Rev. D 23, 165-180 (1981)

[11] V. Silveira and A. Zee, "Scalar phantoms," Phys. Lett. Vol. 161B, No. 1,2,3 (1985) 136-140

[12] D. E. Holz and A. Zee, "Collisional Dark Matter and Scalar Phantoms," arxiv:hep-ph/0105284v1

[13] O. Adriani et al, "Observation of an anomalous positron abundance in the cosmic radiation," Nature 458:607-609,2009 (arXiv:0810.4995v1 [astro-ph])

[14] M. Ackermann et al, "Measurement of separate cosmic-ray electron and positron spectra with the Fermi Large Area Telescope," PRL 108, 011103 (2012) (arXiv:1109.0521v2 [astro-ph.HE])

[15] M. Cirelli, E. Moulin, P. Panci, P. D. Serpico, A. Viana, "Gamma ray constraints on Decaying Dark Matter," arXiv:1205.5283v1

[16] S. F. King, "Vacuum misalignment corrections to tri-bimaximal mixing and form dominance," JHEP 1101:115,2011 (arXiv:1011.6167v2 [hep-ph]); W. Chao and Y. J. Zheng, "Relatively Large Theta13 from Modification to the Tri-bimaximal, Bimaximal and Democratic Neutrino Mixing Matrices," arXiv:1107.0738v1 [hep-ph]


[17] A. Zee, "Obtaining the Neutrino Mixing Matrix with the Tetrahedral Group," Phys.Lett.B630:58-67,2005 (arXiv:hep-ph/0508278v3)

[18] E. Ma, "Neutrino Tribimaximal Mixing from A_4 alone," UCRHEP-T472 (August 2009) (arXiv:0908.3165v2 [hep-ph])

[19] T. P. Cheng, E. Eichten, and L. F. Li, "Higgs phenomena in asymptotically free gauge theories," Phys. Rev. D 9, 22592273 (1974)


[20] K. Nakamura et al. (Particle Data Group), J. Phys. G 37, 075021 (2010) (http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf)

[21] C. T. Hill, C. N. Leung, and S. Rao, "Renormalization group fixed points and the Higgs boson spectrum," Nucl. Phys. B262 (1985) 517-537



(a) $g(\mu)$ (b) $y(\mu)$

Figure 1: The running gauge coupling $g(\mu)$ and Yukawa coupling $y(\mu)$ in the toy model of Eq. (C.4).



(a) $\lambda_1(\mu)$ (b) $\lambda_2(\mu)$

Figure 2: The running quartic self-couplings $\lambda_1(\mu)$ and $\lambda_2(\mu)$ in the toy model of Eq. (C.4).




Figure 3: The running quartic coupling $\lambda_{12}(\mu)$ which couples ϕ_1 and ϕ_2 in the toy model of Eq. (C.4).